

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US04/042981

International filing date: 17 December 2004 (17.12.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/530,828
Filing date: 17 December 2003 (17.12.2003)

Date of receipt at the International Bureau: 24 January 2005 (24.01.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

January 14, 2005

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/530,828

FILING DATE: *December 17, 2003*

RELATED PCT APPLICATION NUMBER: *PCT/US04/42981*



Certified By

Jon W Dudas

Under Secretary
of Commerce for Intellectual Property
and Acting Director of the
United States Patent and Trademark Office

22783 U.S. PTO
121703

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53 (c).

Express Mail Label No. EV342494424US

INVENTOR(S)					
Given Name (first and middle (if any))		Family Name or Surname		Residence (City and either State or Foreign Country)	
Jonathan		Lightner		Mulino, Oregon	
Stephanie K.		Clendennen		Portland, Oregon	
<input type="checkbox"/> Additional inventors are being named on the _____ separately numbered sheets attached hereto					
TITLE OF THE INVENTION (500 characters max)					
GENERATION OF PLANTS WITH ALTERED OIL CONTENT					
CORRESPONDENCE ADDRESS					
Direct all correspondence to:					
<input checked="" type="checkbox"/> Customer Number		23500		Place Customer Number Bar Code Label here	
OR					
<input type="checkbox"/> Firm or Individual Name					
Address					
Address					
City		State		ZIP	
Country		Telephone		Fax	
ENCLOSED APPLICATION PARTS (check all that apply)					
<input checked="" type="checkbox"/> Specification Number of Pages		28		<input type="checkbox"/> CD(s), Number	
<input type="checkbox"/> Drawing(s) Number of Sheets				<input checked="" type="checkbox"/> Other (specify) return receipt postcard	
<input type="checkbox"/> Application Data Sheet. See 37 CFR 1.76					
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT					
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27.					
<input type="checkbox"/> A check or money order is enclosed to cover the filing fees					
<input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number:				50-1108	
<input type="checkbox"/> Payment by credit card. Form PTO-2038 is attached.				FILING FEE AMOUNT (\$) 160	
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.					
<input checked="" type="checkbox"/> No.					
<input type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are: _____.					

Respectfully submitted,
SIGNATURE Jan P. Brunelle
TYPED or PRINTED NAME Jan P. Brunelle
TELEPHONE 650.837.8180

Date 12/17/03
REGISTRATION NO. 35,081
(if appropriate)
Docket Number: AG03-078P

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

This collection of information is required by 37 CFR 1.51. The information is used by the public to file (and by the PTO to process) a provisional application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the complete provisional application to the PTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, Washington, D.C., 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Box Provisional Application, Assistant Commissioner for Patents, Washington, D.C. 20231.

PTO
60/530828
17548
121703

GENERATION OF PLANTS WITH ALTERED OIL CONTENT

BACKGROUND OF THE INVENTION

The ability to manipulate the composition of crop seeds, particularly the content and composition of seed oils, has important applications in the agricultural industries, relating both to processed food oils and to oils for animal feeding. Seeds of agricultural crops contain a variety of valuable constituents, including oil, protein and starch. Industrial processing can separate some or all of these constituents for individual sale in specific applications. For instance, nearly 60% of the US soybean crop is crushed by the soy processing industry. Soy processing yields purified oil, which is sold at high value, while the remainder is sold principally for lower value livestock feed (US Soybean Board, 2001 Soy Stats). Canola seed is crushed to produce oil and the co-product canola meal (Canola Council of Canada). Nearly 20% of the 1999/2000 US corn crop was industrially refined, primarily for production of starch, ethanol and oil (Corn Refiners Association). Thus, it is often desirable to maximize oil content of seeds. For instance, for processed oilseeds such as soy and canola, increasing the absolute oil content of the seed will increase the value of such grains. For processed corn it may be desired to either increase or decrease oil content, depending on utilization of other major constituents. Decreasing oil may improve the quality of isolated starch by reducing undesired flavors associated with oil oxidation. Alternatively, in ethanol production, where flavor is unimportant, increasing oil content may increase overall value. In many fed grains, such as corn and wheat, it is desirable to increase seed oil content, because oil has higher energy content than other seed constituents such as carbohydrate. Oilseed processing, like most grain processing businesses, is a capital-intensive business; thus small shifts in the distribution of products from the low valued components to the high value oil component can have substantial economic impacts for grain processors.

Biotechnological manipulation of oils can provide compositional alteration and improvement of oil yield. Compositional alterations include high oleic soybean and corn oil (US Pat Nos 6,229,033 and 6,248,939), and laurate-containing seeds (US Pat No 5,639,790), among others. Work in compositional alteration has predominantly focused on processed oilseeds but has been readily extendable to non-oilseed crops, including corn. While there is considerable interest in increasing oil content, the only currently practiced biotechnology in this area is High-Oil Corn (HOC) technology (DuPont, US PAT NO: 5,704,160). HOC employs high oil pollinators developed by classical selection breeding along with elite

(male-sterile) hybrid females in a production system referred to as TopCross. The TopCross High Oil system raises harvested grain oil content in maize from ~3.5% to ~7%, improving the energy content of the grain.

While it has been fruitful, the HOC production system has inherent limitations.

5 First, the system of having a low percentage of pollinators responsible for an entire field's seed set contains inherent risks, particularly in drought years. Second, oil contents in current HOC fields have plateaued at about 9% oil. Finally, high-oil corn is not primarily a biochemical change, but rather an anatomical mutant (increased embryo size) that has the indirect result of increasing oil content. For these reasons, an alternative high oil strategy,
10 particularly one that derives from an altered biochemical output, would be especially valuable.

The most obvious target crops for the processed oil market are soy and rapeseed, and a large body of commercial work (e.g., US Pat No: 5,952,544; PCT application WO9411516) demonstrates that *Arabidopsis* is an excellent model for oil metabolism in
15 these crops. Biochemical screens of seed oil composition have identified *Arabidopsis* genes for many critical biosynthetic enzymes and have led to identification of agronomically important gene orthologs. For instance, screens using chemically mutagenized populations have identified lipid mutants whose seeds display altered fatty acid composition (Lemieux et al., 1990; James and Dooner, 1990). T-DNA mutagenesis screens (Feldmann et al., 1989)
20 that detected altered fatty acid composition identified the omega 3 desaturase (*FAD3*) and delta-12 desaturase (*FAD2*) genes (US Pat No 5952544; Yadav et al., 1993; Okuley et al., 1994). A screen which focused on oil content rather than oil quality, analyzed chemically-induced mutants for wrinkled seeds or altered seed density, from which altered seed oil content was inferred (Focks and Benning, 1998). Another screen, designed to identify
25 enzymes involved in production of very long chain fatty acids, identified a mutation in the gene encoding a diacylglycerol acyltransferase (DGAT) as being responsible for reduced triacyl glycerol accumulation in seeds (Katavic V et al, 1995). It was further shown that seed-specific over-expression of the DGAT cDNA was associated with increased seed oil content (Jako et al., 2001).

30 Activation tagging in plants refers to a method of generating random mutations by insertion of a heterologous nucleic acid construct comprising regulatory sequences (e.g., an enhancer) into a plant genome. The regulatory sequences can act to enhance transcription of one or more native plant genes; accordingly, activation tagging is a fruitful method for generating gain-of-function, generally dominant mutants (see, e.g., Hayashi *et al.*, 1992;

ability to incorporate and express heterologous DNA fragments in a foreign cell. Many prokaryotic and eukaryotic expression vectors are commercially available. Selection of appropriate expression vectors is within the knowledge of those having skill in the art.

5 A "heterologous" nucleic acid construct or sequence has a portion of the sequence that is not native to the plant cell in which it is expressed. Heterologous, with respect to a control sequence refers to a control sequence (*i.e.* promoter or enhancer) that does not function in nature to regulate the same gene the expression of which it is currently regulating. Generally, heterologous nucleic acid sequences are not endogenous to the cell or part of the genome in which they are present, and have been added to the cell, by infection, 10 transfection, microinjection, electroporation, or the like. A "heterologous" nucleic acid construct may contain a control sequence/DNA coding sequence combination that is the same as, or different from a control sequence/DNA coding sequence combination found in the native plant.

15 As used herein, the term "gene" means the segment of DNA involved in producing a polypeptide chain, which may or may not include regions preceding and following the coding region, *e.g.* 5' untranslated (5' UTR) or "leader" sequences and 3' UTR or "trailer" sequences, as well as intervening sequences (introns) between individual coding segments (exons) and non-transcribed regulatory sequence.

20 As used herein, "recombinant" includes reference to a cell or vector, that has been modified by the introduction of a heterologous nucleic acid sequence or that the cell is derived from a cell so modified. Thus, for example, recombinant cells express genes that are not found in identical form within the native (non-recombinant) form of the cell or express native genes that are otherwise abnormally expressed, under expressed or not expressed at all as a result of deliberate human intervention.

25 As used herein, the term "gene expression" refers to the process by which a polypeptide is produced based on the nucleic acid sequence of a gene. The process includes both transcription and translation; accordingly, "expression" may refer to either a polynucleotide or polypeptide sequence, or both. Sometimes, expression of a polynucleotide sequence will not lead to protein translation. "Over-expression" refers to 30 increased expression of a polynucleotide and/or polypeptide sequence relative to its expression in a wild-type (or other reference [*e.g.*, non-transgenic]) plant and may relate to a naturally-occurring or non-naturally occurring sequence. "Ectopic expression" refers to expression at a time, place, and/or increased level that does not naturally occur in the non-altered or wild-type plant. "Under-expression" refers to decreased expression of a

polynucleotide and/or polypeptide sequence, generally of an endogenous gene, relative to its expression in a wild-type plant. The terms "mis-expression" and "altered expression" encompass over-expression, under-expression, and ectopic expression.

The term "introduced" in the context of inserting a nucleic acid sequence into a cell, means "transfection", or "transformation" or "transduction" and includes reference to the incorporation of a nucleic acid sequence into a eukaryotic or prokaryotic cell where the nucleic acid sequence may be incorporated into the genome of the cell (for example, chromosome, plasmid, plastid, or mitochondrial DNA), converted into an autonomous replicon, or transiently expressed (for example, transfected mRNA).

As used herein, a "plant cell" refers to any cell derived from a plant, including cells from undifferentiated tissue (*e.g.*, callus) as well as plant seeds, pollen, progagules and embryos.

As used herein, the terms "native" and "wild-type" relative to a given plant trait or phenotype refers to the form in which that trait or phenotype is found in the same variety of plant in nature.

As used herein, the term "modified" regarding a plant trait, refers to a change in the phenotype of a transgenic plant relative to the similar non-transgenic plant. An "interesting phenotype (trait)" with reference to a transgenic plant refers to an observable or measurable phenotype demonstrated by a T1 and/or subsequent generation plant, which is not displayed by the corresponding non-transgenic (*i.e.*, a genotypically similar plant that has been raised or assayed under similar conditions). An interesting phenotype may represent an improvement in the plant or may provide a means to produce improvements in other plants. An "improvement" is a feature that may enhance the utility of a plant species or variety by providing the plant with a unique and/or novel quality. An "altered oil content phenotype" refers to measurable phenotype of a genetically modified plant, where the plant displays a statistically significant increase or decrease in overall oil content (*i.e.*, the percentage of seed mass that is oil), as compared to the similar, but non-modified plant. A high oil phenotype refers to an increase in overall oil content.

As used herein, a "mutant" polynucleotide sequence or gene differs from the corresponding wild type polynucleotide sequence or gene either in terms of sequence or expression, where the difference contributes to a modified plant phenotype or trait. Relative to a plant or plant line, the term "mutant" refers to a plant or plant line which has a modified plant phenotype or trait, where the modified phenotype or trait is associated with the modified expression of a wild type polynucleotide sequence or gene.

As used herein, the term "T1" refers to the generation of plants from the seed of T0 plants. The T1 generation is the first set of transformed plants that can be selected by application of a selection agent, *e.g.*, an antibiotic or herbicide, for which the transgenic plant contains the corresponding resistance gene. The term "T2" refers to the generation of plants by self-fertilization of the flowers of T1 plants, previously selected as being transgenic. T3 plants are generated from T2 plants, etc. As used herein, the "direct progeny" of a given plant derives from the seed (or, sometimes, other tissue) of that plant and is in the immediately subsequent generation; for instance, for a given lineage, a T2 plant is the direct progeny of a T1 plant. The "indirect progeny" of a given plant derives from the seed (or other tissue) of the direct progeny of that plant, or from the seed (or other tissue) of subsequent generations in that lineage; for instance, a T3 plant is the indirect progeny of a T1 plant.

As used herein, the term "plant part" includes any plant organ or tissue, including, without limitation, seeds, embryos, meristematic regions, callus tissue, leaves, roots, shoots, gametophytes, sporophytes, pollen, and microspores. Plant cells can be obtained from any plant organ or tissue and cultures prepared therefrom. The class of plants which can be used in the methods of the present invention is generally as broad as the class of higher plants amenable to transformation techniques, including both monocotyledenous and dicotyledenous plants.

As used herein, "transgenic plant" includes a plant that comprises within its genome a heterologous polynucleotide. The heterologous polynucleotide can be either stably integrated into the genome, or can be extra-chromosomal. Preferably, the polynucleotide of the present invention is stably integrated into the genome such that the polynucleotide is passed on to successive generations. A plant cell, tissue, organ, or plant into which the heterologous polynucleotides have been introduced is considered "transformed", "transfected", or "transgenic". Direct and indirect progeny of transformed plants or plant cells that also contain the heterologous polynucleotide are also considered transgenic.

Identification of Plants with an Altered Oil Content Phenotype

We used an *Arabidopsis* activation tagging screen to identify the association between the gene we have designated "HIO30.5," (At3g52280; GI#18409525:1-1110), and an altered oil content phenotype (specifically, a high oil phenotype). Briefly, and as further described in the Examples, a large number of *Arabidopsis* plants were mutated with the pSKI015 vector, which comprises a T-DNA from the Ti plasmid of *Agrobacterium tumefaciens*, a

viral enhancer element, and a selectable marker gene (Weigel *et al*, 2000). When the T-DNA inserts into the genome of transformed plants, the enhancer element can cause up-regulation genes in the vicinity, generally within about 10 kilobase (kb) of the insertion. T1 plants were exposed to the selective agent in order to specifically recover transformed plants that expressed the selectable marker and therefore harbored T-DNA insertions. Samples of approximately 15-20 T2 seeds were collected from transformed T1 plants, and lipids were extracted from whole seeds. Gas chromatography (GC) analysis was performed to determine fatty acid content and composition of seed samples.

An *Arabidopsis* line that showed a high-oil phenotype was identified, wherein oils (i.e., fatty acids) constituted approximately 35% of seed mass. The association of the HIO30.5 gene with the high oil phenotype was discovered by analysis of the genomic DNA sequence flanking the T-DNA insertion in the identified line. Accordingly, HIO30.5 genes and/or polypeptides may be employed in the development of genetically modified plants having a modified oil content phenotype ("a HIO30.5 phenotype"). HIO30.5 genes may be used in the generation of oilseed crops that provide improved oil yield from oilseed processing and in the generation of feed grain crops that provide increased energy for animal feeding. HIO30.5 genes may further be used to increase the oil content of specialty oil crops, in order to augment yield of desired unusual fatty acids. Transgenic plants that have been genetically modified to express HIO30.5 can be used in the production of oil, wherein the transgenic plants are grown, and oil is obtained from plant parts (e.g. seed) using standard methods.

HIO30.5 Nucleic Acids and Polypeptides

Arabidopsis HIO30.5 nucleic acid (genomic DNA) sequence is provided in SEQ ID NO:1 and in Genbank entry GI#18409525:1-1110. The corresponding protein sequence is provided in SEQ ID NO:2 and in GI#15231174. Nucleic acids and/or proteins that are orthologs or paralogs of *Arabidopsis* HIO30.5, are described in Example 3 below.

As used herein, the term "HIO30.5 polypeptide" refers to a full-length HIO30.5 protein or a fragment, derivative (variant), or ortholog thereof that is "functionally active," meaning that the protein fragment, derivative, or ortholog exhibits one or more of the functional activities associated with the polypeptide of SEQ ID NO:2. In one preferred embodiment, a functionally active HIO30.5 polypeptide causes an altered oil content phenotype when mis-expressed in a plant. In a further preferred embodiment, mis-expression of the HIO30.5 polypeptide causes a high oil phenotype in a plant. In another

embodiment, a functionally active HIO30.5 polypeptide is capable of rescuing defective (including deficient) endogenous HIO30.5 activity when expressed in a plant or in plant cells; the rescuing polypeptide may be from the same or from a different species as that with defective activity. In another embodiment, a functionally active fragment of a full length HIO30.5 polypeptide (i.e., a native polypeptide having the sequence of SEQ ID NO:2 or a naturally occurring ortholog thereof) retains one or more of the biological properties associated with the full-length HIO30.5 polypeptide, such as signaling activity, binding activity, catalytic activity, or cellular or extra-cellular localizing activity. A HIO30.5 fragment preferably comprises a beta subunit of transcription initiation factor IIF domain HIO30.5 domain, such as a C- or N-terminal or catalytic domain, among others, and preferably comprises at least 10, preferably at least 20, more preferably at least 25, and most preferably at least 50 contiguous amino acids of a HIO30.5 protein. Functional domains can be identified using the PFAM program (Bateman A et al., 1999 Nucleic Acids Res 27:260-262; website at pfam.wustl.edu). A preferred HIO30.5 fragment comprises a bromodomain (PFam00439). Functionally active variants of full-length HIO30.5 polypeptides or fragments thereof include polypeptides with amino acid insertions, deletions, or substitutions that retain one or more of the biological properties associated with the full-length HIO30.5 polypeptide. In some cases, variants are generated that change the post-translational processing of a HIO30.5 polypeptide. For instance, variants may have altered protein transport or protein localization characteristics or altered protein half-life compared to the native polypeptide.

As used herein, the term "HIO30.5 nucleic acid" encompasses nucleic acids with the sequence provided in or complementary to the sequence provided in SEQ ID NO:1, as well as functionally active fragments, derivatives, or orthologs thereof. A HIO30.5 nucleic acid of this invention may be DNA, derived from genomic DNA or cDNA, or RNA.

In one embodiment, a functionally active HIO30.5 nucleic acid encodes or is complementary to a nucleic acid that encodes a functionally active HIO30.5 polypeptide. Included within this definition is genomic DNA that serves as a template for a primary RNA transcript (i.e., an mRNA precursor) that requires processing, such as splicing, before encoding the functionally active HIO30.5 polypeptide. A HIO30.5 nucleic acid can include other non-coding sequences, which may or may not be transcribed; such sequences include 5' and 3' UTRs, polyadenylation signals and regulatory sequences that control gene expression, among others, as are known in the art. Some polypeptides require processing events, such as proteolytic cleavage, covalent modification, etc., in order to become fully

active. Accordingly, functionally active nucleic acids may encode the mature or the pre-processed HIO30.5 polypeptide, or an intermediate form. A HIO30.5 polynucleotide can also include heterologous coding sequences, for example, sequences that encode a marker included to facilitate the purification of the fused polypeptide, or a transformation marker.

5 In another embodiment, a functionally active HIO30.5 nucleic acid is capable of being used in the generation of loss-of-function HIO30.5 phenotypes, for instance, via antisense suppression, co-suppression, etc.

In one preferred embodiment, a HIO30.5 nucleic acid used in the methods of this invention comprises a nucleic acid sequence that encodes or is complementary to a sequence
10 that encodes a HIO30.5 polypeptide having at least 50%, 60%, 70%, 75%, 80%, 85%, 90%, 95% or more sequence identity to the polypeptide sequence presented in SEQ ID NO:2.

In another embodiment a HIO30.5 polypeptide of the invention comprises a polypeptide sequence with at least 50% or 60% identity to the HIO30.5 polypeptide sequence of SEQ ID NO:2, and may have at least 70%, 80%, 85%, 90% or 95% or more
15 sequence identity to the HIO30.5 polypeptide sequence of SEQ ID NO:2. In another embodiment, a HIO30.5 polypeptide comprises a polypeptide sequence with at least 50%, 60%, 70%, 80%, 85%, 90% or 95% or more sequence identity to a functionally active fragment of the polypeptide presented in SEQ ID NO:2, such as a bromodomain. In yet another embodiment, a HIO30.5 polypeptide comprises a polypeptide sequence with at least
20 50%, 60 %, 70%, 80%, or 90% identity to the polypeptide sequence of SEQ ID NO:2 over its entire length and comprises a bromodomain.

In another aspect, a HIO30.5 polynucleotide sequence is at least 50% to 60% identical over its entire length to the HIO30.5 nucleic acid sequence presented as SEQ ID NO:1, or nucleic acid sequences that are complementary to such a HIO30.5 sequence, and
25 may comprise at least 70%, 80%, 85%, 90% or 95% or more sequence identity to the HIO30.5 sequence presented as SEQ ID NO:1 or a functionally active fragment thereof, or complementary sequences.

As used herein, "percent (%) sequence identity" with respect to a specified subject sequence, or a specified portion thereof, is defined as the percentage of nucleotides or amino
30 acids in the candidate derivative sequence identical with the nucleotides or amino acids in the subject sequence (or specified portion thereof), after aligning the sequences and introducing gaps, if necessary to achieve the maximum percent sequence identity, as generated by the program WU-BLAST-2.0a19 (Altschul *et al.*, J. Mol. Biol. (1997) 215:403-410; website at blast.wustl.edu/blast/README.html) with search parameters set to

default values. The HSP S and HSP S2 parameters are dynamic values and are established by the program itself depending upon the composition of the particular sequence and composition of the particular database against which the sequence of interest is being searched. A "% identity value" is determined by the number of matching identical nucleotides or amino acids divided by the sequence length for which the percent identity is being reported. "Percent (%) amino acid sequence similarity" is determined by doing the same calculation as for determining % amino acid sequence identity, but including conservative amino acid substitutions in addition to identical amino acids in the computation. A conservative amino acid substitution is one in which an amino acid is substituted for another amino acid having similar properties such that the folding or activity of the protein is not significantly affected. Aromatic amino acids that can be substituted for each other are phenylalanine, tryptophan, and tyrosine; interchangeable hydrophobic amino acids are leucine, isoleucine, methionine, and valine; interchangeable polar amino acids are glutamine and asparagine; interchangeable basic amino acids are arginine, lysine and histidine; interchangeable acidic amino acids are aspartic acid and glutamic acid; and interchangeable small amino acids are alanine, serine, threonine, cysteine and glycine.

Derivative nucleic acid molecules of the subject nucleic acid molecules include sequences that selectively hybridize to the nucleic acid sequence of SEQ ID NO:1. The stringency of hybridization can be controlled by temperature, ionic strength, pH, and the presence of denaturing agents such as formamide during hybridization and washing. Conditions routinely used are well known (see, *e.g.*, Current Protocol in Molecular Biology, Vol. 1, Chap. 2.10, John Wiley & Sons, Publishers (1994); Sambrook *et al.*, Molecular Cloning, Cold Spring Harbor (1989)). In some embodiments, a nucleic acid molecule of the invention is capable of hybridizing to a nucleic acid molecule containing the nucleotide sequence of SEQ ID NO:1 under stringent hybridization conditions that are: prehybridization of filters containing nucleic acid for 8 hours to overnight at 65° C in a solution comprising 6X single strength citrate (SSC) (1X SSC is 0.15 M NaCl, 0.015 M Na citrate; pH 7.0), 5X Denhardt's solution, 0.05% sodium pyrophosphate and 100 µg/ml herring sperm DNA; hybridization for 18-20 hours at 65° C in a solution containing 6X SSC, 1X Denhardt's solution, 100 µg/ml yeast tRNA and 0.05% sodium pyrophosphate; and washing of filters at 65° C for 1 h in a solution containing 0.1X SSC and 0.1% SDS (sodium dodecyl sulfate). In other embodiments, moderately stringent hybridization conditions are used that are: pretreatment of filters containing nucleic acid for 6 h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.1% PVP,

0.1% Ficoll, 1% BSA, and 500 $\mu\text{g/ml}$ denatured salmon sperm DNA; hybridization for 18-20 h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 $\mu\text{g/ml}$ salmon sperm DNA, and 10% (wt/vol) dextran sulfate; followed by washing twice for 1 hour at 55° C in a solution containing 2X SSC and 0.1% SDS. Alternatively, low stringency conditions can be used that comprise: incubation for 8 hours to overnight at 37° C in a solution comprising 20% formamide, 5 x SSC, 50 mM sodium phosphate (pH 7.6), 5X Denhardt's solution, 10% dextran sulfate, and 20 $\mu\text{g/ml}$ denatured sheared salmon sperm DNA; hybridization in the same buffer for 18 to 20 hours; and washing of filters in 1 x SSC at about 37° C for 1 hour.

As a result of the degeneracy of the genetic code, a number of polynucleotide sequences encoding a HIO30.5 polypeptide can be produced. For example, codons may be selected to increase the rate at which expression of the polypeptide occurs in a particular host species, in accordance with the optimum codon usage dictated by the particular host organism (see, e.g., Nakamura et al, 1999). Such sequence variants may be used in the methods of this invention.

The methods of the invention may use orthologs of the *Arabidopsis* HIO30.5. Methods of identifying the orthologs in other plant species are known in the art. Normally, orthologs in different species retain the same function, due to presence of one or more protein motifs and/or 3-dimensional structures. In evolution, when a gene duplication event follows speciation, a single gene in one species, such as *Arabidopsis*, may correspond to multiple genes (paralogs) in another. As used herein, the term "orthologs" encompasses paralogs. When sequence data is available for a particular plant species, orthologs are generally identified by sequence homology analysis, such as BLAST analysis, usually using protein bait sequences. Sequences are assigned as a potential ortholog if the best hit sequence from the forward BLAST result retrieves the original query sequence in the reverse BLAST (Huynen MA and Bork P, Proc Natl Acad Sci (1998) 95:5849-5856; Huynen MA *et al.*, Genome Research (2000) 10:1204-1210). Programs for multiple sequence alignment, such as CLUSTAL (Thompson JD et al, 1994, Nucleic Acids Res 22:4673-4680) may be used to highlight conserved regions and/or residues of orthologous proteins and to generate phylogenetic trees. In a phylogenetic tree representing multiple homologous sequences from diverse species (e.g., retrieved through BLAST analysis), orthologous sequences from two species generally appear closest on the tree with respect to all other sequences from these two species. Structural threading or other analysis of protein folding (e.g., using software by ProCeryon, Biosciences, Salzburg, Austria) may also

identify potential orthologs. Nucleic acid hybridization methods may also be used to find orthologous genes and are preferred when sequence data are not available. Degenerate PCR and screening of cDNA or genomic DNA libraries are common methods for finding related gene sequences and are well known in the art (see, e.g., Sambrook, 1989; Dieffenbach and Dveksler, 1995). For instance, methods for generating a cDNA library from the plant species of interest and probing the library with partially homologous gene probes are described in Sambrook *et al.* A highly conserved portion of the *Arabidopsis* HIO30.5 coding sequence may be used as a probe. HIO30.5 ortholog nucleic acids may hybridize to the nucleic acid of SEQ ID NO:1 under high, moderate, or low stringency conditions. After amplification or isolation of a segment of a putative ortholog, that segment may be cloned and sequenced by standard techniques and utilized as a probe to isolate a complete cDNA or genomic clone. Alternatively, it is possible to initiate an EST project to generate a database of sequence information for the plant species of interest. In another approach, antibodies that specifically bind known HIO30.5 polypeptides are used for ortholog isolation (see, e.g., Harlow and Lane, 1988, 1999). Western blot analysis can determine that a HIO30.5 ortholog (i.e., an orthologous protein) is present in a crude extract of a particular plant species. When reactivity is observed, the sequence encoding the candidate ortholog may be isolated by screening expression libraries representing the particular plant species. Expression libraries can be constructed in a variety of commercially available vectors, including lambda gt11, as described in Sambrook, *et al.*, 1989. Once the candidate ortholog(s) are identified by any of these means, candidate orthologous sequence are used as bait (the "query") for the reverse BLAST against sequences from *Arabidopsis* or other species in which HIO30.5 nucleic acid and/or polypeptide sequences have been identified.

HIO30.5 nucleic acids and polypeptides may be obtained using any available method. For instance, techniques for isolating cDNA or genomic DNA sequences of interest by screening DNA libraries or by using polymerase chain reaction (PCR), as previously described, are well known in the art. Alternatively, nucleic acid sequence may be synthesized. Any known method, such as site directed mutagenesis (Kunkel TA *et al.*, 1991), may be used to introduce desired changes into a cloned nucleic acid.

In general, the methods of the invention involve incorporating the desired form of the HIO30.5 nucleic acid into a plant expression vector for transformation of in plant cells, and the HIO30.5 polypeptide is expressed in the host plant.

An isolated HIO30.5 nucleic acid molecule is other than in the form or setting in which it is found in nature and is identified and separated from least one contaminant

nucleic acid molecule with which it is ordinarily associated in the natural source of the HIO30.5 nucleic acid. However, an isolated HIO30.5 nucleic acid molecule includes HIO30.5 nucleic acid molecules contained in cells that ordinarily express HIO30.5 where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

Generation of Genetically Modified Plants with an Altered Oil Content

Phenotype

HIO30.5 nucleic acids and polypeptides may be used in the generation of genetically modified plants having a modified oil content phenotype. As used herein, a "modified oil content phenotype" may refer to modified oil content in any part of the plant; the modified oil content is often observed in seeds. In a preferred embodiment, altered expression of the HIO30.5 gene in a plant is used to generate plants with a high oil phenotype.

The methods described herein are generally applicable to all plants. Although activation tagging and gene identification is carried out in *Arabidopsis*, the HIO30.5 gene (or an ortholog, variant or fragment thereof) may be expressed in any type of plant. In a preferred embodiment, the invention is directed to oil-producing plants, which produce and store triacylglycerol in specific organs, primarily in seeds. Such species include soybean (*Glycine max*), rapeseed and canola (including *Brassica napus*, *B. campestris*), sunflower (*Helianthus annuus*), cotton (*Gossypium hirsutum*), corn (*Zea mays*), cocoa (*Theobroma cacao*), safflower (*Carthamus tinctorius*), oil palm (*Elaeis guineensis*), coconut palm (*Cocos nucifera*), flax (*Linum usitatissimum*), castor (*Ricinus communis*) and peanut (*Arachis hypogaea*). The invention may also be directed to fruit- and vegetable-bearing plants, grain-producing plants, nut-producing plants, rapid cycling *Brassica* species, alfalfa (*Medicago sativa*), tobacco (*Nicotiana*), turfgrass (*Poaceae* family), other forage crops, and wild species that may be a source of unique fatty acids.

The skilled artisan will recognize that a wide variety of transformation techniques exist in the art, and new techniques are continually becoming available. Any technique that is suitable for the target host plant can be employed within the scope of the present invention. For example, the constructs can be introduced in a variety of forms including, but not limited to as a strand of DNA, in a plasmid, or in an artificial chromosome. The introduction of the constructs into the target plant cells can be accomplished by a variety of techniques, including, but not limited to *Agrobacterium*-mediated transformation, electroporation, microinjection, microprojectile bombardment calcium-phosphate-DNA co-

precipitation or liposome-mediated transformation of a heterologous nucleic acid. The transformation of the plant is preferably permanent, *i.e.* by integration of the introduced expression constructs into the host plant genome, so that the introduced constructs are passed onto successive plant generations. Depending upon the intended use, a heterologous nucleic acid construct comprising an HIO30.5 polynucleotide may encode the entire protein or a biologically active portion thereof.

In one embodiment, binary Ti-based vector systems may be used to transfer polynucleotides. Standard *Agrobacterium* binary vectors are known to those of skill in the art, and many are commercially available (e.g., pBI121 Clontech Laboratories, Palo Alto, CA).

The optimal procedure for transformation of plants with *Agrobacterium* vectors will vary with the type of plant being transformed. Exemplary methods for *Agrobacterium*-mediated transformation include transformation of explants of hypocotyl, shoot tip, stem or leaf tissue, derived from sterile seedlings and/or plantlets. Such transformed plants may be reproduced sexually, or by cell or tissue culture. *Agrobacterium* transformation has been previously described for a large number of different types of plants and methods for such transformation may be found in the scientific literature. Of particular relevance are methods to transform commercially important crops, such as rapeseed (De Block et al., 1989), sunflower (Everett et al., 1987), and soybean (Christou et al., 1989; Kline et al., 1987).

Expression (including transcription and translation) of HIO30.5 may be regulated with respect to the level of expression, the tissue type(s) where expression takes place and/or developmental stage of expression. A number of heterologous regulatory sequences (e.g., promoters and enhancers) are available for controlling the expression of a HIO30.5 nucleic acid. These include constitutive, inducible and regulatable promoters, as well as promoters and enhancers that control expression in a tissue- or temporal-specific manner. Exemplary constitutive promoters include the raspberry E4 promoter (U.S. Patent Nos. 5,783,393 and 5,783,394), the 35S CaMV (Jones JD *et al.*, 1992), the CsVMV promoter (Verdaguer B *et al.*, 1998) and the melon actin promoter (published PCT application WO0056863). Exemplary tissue-specific promoters include the tomato E4 and E8 promoters (U.S. Patent No. 5,859,330) and the tomato 2AII gene promoter (Van Haaren MJJ *et al.*, 1993).

In one preferred embodiment, HIO30.5 expression is under control of regulatory sequences from genes whose expression is associated with early seed and/or embryo development. Legume genes whose promoters are associated with early seed and embryo development include *V. faba legumin* (Baumlein et al., 1991, Mol Gen Genet 225:121-8;

Baumlein et al., 1992, Plant J 2:233-9), *V. faba usp* (Fiedler et al., 1993, Plant Mol Biol 22:669-79), pea *convicilin* (Bown et al., 1988, Biochem J 251:717-26), pea *lectin* (dePater et al., 1993, Plant Cell 5:877-86), *P. vulgaris beta phaseolin* (Bustos et al., 1991, EMBO J 10:1469-79), *P. vulgaris DLEC2* and *PHS [beta]* (Bobb et al, 1997, Nucleic Acids Res 25:641-7), and soybean *beta-Conglycinin*, 7S storage protein (Chamberland et al., 1992, Plant Mol Biol 19:937-49). Cereal genes whose promoters are associated with early seed and embryo development include rice *glutelin* ("GluA-3," Yoshihara and Takaiwa, 1996, Plant Cell Physiol 37:107-11; "GluB-1," Takaiwa et al., 1996, Plant Mol Biol 30:1207-21; Washida et al., 1999, Plant Mol Biol 40:1-12; "Gt3," Leisy et al., 1990, Plant Mol Biol 14:41-50), rice *prolamin* (Zhou & Fan, 1993, Transgenic Res 2:141-6), wheat *prolamin* (Hammond-Kosack et al., 1993, EMBO J 12:545-54), maize *zein* (Z4, Matzke et al., 1990, Plant Mol Biol 14:323-32), and barley *B-hordeins* (Entwistle et al., 1991, Plant Mol Biol 17:1217-31). Other genes whose promoters are associated with early seed and embryo development include oil palm GLO7A (7S globulin, Morcillo et al., 2001, Physiol Plant 112:233-243), *Brassica napus napin*, 2S storage protein, and napA gene (Josefsson et al., 1987, J Biol Chem 262:12196-201; Stalberg et al., 1993, Plant Mol Biol 1993 23:671-83; Ellerstrom et al., 1996, Plant Mol Biol 32:1019-27), *Brassica napus oleosin* (Keddie et al., 1994, Plant Mol Biol 24:327-40), *Arabidopsis oleosin* (Plant et al., 1994, Plant Mol Biol 25:193-205), *Arabidopsis* FAE1 (Rossak et al., 2001, Plant Mol Biol 46:717-25), *Canavalia gladiata* conA (Yamamoto et al., 1995, Plant Mol Biol 27:729-41), and *Catharanthus roseus* strictosidine synthase (Str, Ouwerkerk and Memelink, 1999, Mol Gen Genet 261:635-43). In another preferred embodiment, regulatory sequences from genes expressed during oil biosynthesis are used (see, e.g., US Pat No: 5,952, 544). Alternative promoters are from plant storage protein genes (Bevan et al, 1993, Philos Trans R Soc Lond B Biol Sci 342:209-15).

In yet another aspect, in some cases it may be desirable to inhibit the expression of endogenous HIO30.5 in a host cell. Exemplary methods for practicing this aspect of the invention include, but are not limited to antisense suppression (Smith, *et al.*, 1988; van der Krol et al., 1988); co-suppression (Napoli, *et al.*, 1990); ribozymes (PCT Publication WO 97/10328); and combinations of sense and antisense (Waterhouse, *et al.*, 1998). Methods for the suppression of endogenous sequences in a host cell typically employ the transcription or transcription and translation of at least a portion of the sequence to be suppressed. Such sequences may be homologous to coding as well as non-coding regions of the endogenous sequence. Antisense inhibition may use the entire cDNA sequence (Sheehy et al., 1988), a

partial cDNA sequence including fragments of 5' coding sequence, (Cannon et al., 1990), or 3' non-coding sequences (Ch'ng et al., 1989). Cosuppression techniques may use the entire cDNA sequence (Napoli et al., 1990; van der Krol et al., 1990), or a partial cDNA sequence (Smith et al., (1990).

5

Standard molecular and genetic tests may be performed to further analyze the association between a gene and an observed phenotype. Exemplary techniques are described below.

1. DNA/RNA analysis

10

The stage- and tissue-specific gene expression patterns in mutant versus wild-type lines may be determined, for instance, by in situ hybridization. Analysis of the methylation status of the gene, especially flanking regulatory regions, may be performed. Other suitable techniques include overexpression, ectopic expression, expression in other plant species and gene knock-out (reverse genetics, targeted knock-out, viral induced gene silencing [VIGS, see Baulcombe D, 1999]).

15

In a preferred application expression profiling, generally by microarray analysis, is used to simultaneously measure differences or induced changes in the expression of many different genes. Techniques for microarray analysis are well known in the art (Schena M *et al.*, Science (1995) 270:467-470; Baldwin D *et al.*, 1999; Dangond F, Physiol Genomics (2000) 2:53-58; van Hal NL *et al.*, J Biotechnol (2000) 78:271-280; Richmond T and Somerville S, Curr Opin Plant Biol (2000) 3:108-116). Expression profiling of individual tagged lines may be performed. Such analysis can identify other genes that are coordinately regulated as a consequence of the overexpression of the gene of interest, which may help to place an unknown gene in a particular pathway.

20

2. Gene Product Analysis

25

Analysis of gene products may include recombinant protein expression, antisera production, immunolocalization, biochemical assays for catalytic or other activity, analysis of phosphorylation status, and analysis of interaction with other proteins via yeast two-hybrid assays.

30

3. Pathway Analysis

Pathway analysis may include placing a gene or gene product within a particular biochemical, metabolic or signaling pathway based on its mis-expression phenotype or by sequence homology with related genes. Alternatively, analysis may comprise genetic crosses with wild-type lines and other mutant lines (creating double mutants) to order the

gene in a pathway, or determining the effect of a mutation on expression of downstream "reporter" genes in a pathway.

While the invention has been described with reference to specific methods and
embodiments, it will be appreciated that various modifications and changes may be made
without departing from the invention. All publications cited herein are expressly
incorporated herein by reference for the purpose of describing and disclosing compositions
and methodologies that might be used in connection with the invention. All cited patents,
patent applications, and sequence information in referenced websites and public databases
are also incorporated by reference.

EXAMPLES

EXAMPLE 1

Generation of Plants with a HIO30.5 Phenotype by Transformation with an Activation Tagging Construct

Mutants were generated using the activation tagging "ACTTAG" vector, pSKI015
(GI#6537289; Weigel D *et al.*, 2000). Standard methods were used for the generation of
Arabidopsis transgenic plants, and were essentially as described in published application
PCT WO0183697. Briefly, T0 *Arabidopsis* (Col-0) plants were transformed with
Agrobacterium carrying the pSKI015 vector, which comprises T-DNA derived from the
Agrobacterium Ti plasmid, an herbicide resistance selectable marker gene, and the 4X
CaMV 35S enhancer element. Transgenic plants were selected at the T1 generation based
on herbicide resistance. T2 seed was collected from T1 plants and stored in an indexed
collection, and a portion of the T2 seed was accessed for the screen.

Quantitative determination of seed fatty acid content was performed using the
follows methods. An aliquot of 15 to 20 T2 seeds from each line tested, which generally
contained homozygous insertion, homozygous wild-type, and heterozygous genotypes in a
standard 1:1:2 ratio, was massed on UMT-2 ultra-microbalance (Mettler-Toledo Co., Ohio,
USA) and then transferred to a glass extraction vial. Whole seeds were trans-esterified in
500 ul 2.5% H₂SO₄ in MeOH for 3 hours at 80 °C, following the method of Browse *et al.*
(Biochem J 235:25-31, 1986) with modifications. A known amount of heptadecanoic acid
was included in the reaction as an internal standard. 750 ul of water and 400 ul of hexane

were added to each vial, which was then shaken vigorously and allowed to phase separate. Reaction vials were loaded directly onto GC for analysis and the upper hexane phase was sampled by the autosampler. Gas chromatography with Flame Ionization detection was used to separate and quantify the fatty acid methyl esters. Agilent 6890 Plus GC's were used for separation with Agilent Innowax columns (30m x 0.25mm ID, 250um film thickness). The carrier gas was Hydrogen at a constant flow of 2.5 ml/ minute. 1ul of sample was injected in splitless mode (inlet temperature 220°C, Purge flow 15ml/min at 1 minute). The oven was programmed for an initial temperature of 105°C, initial time 0.5 minutes, followed by a ramp of 60°C per minute to 175°C, a 40°C /minute ramp to 260°C with a final hold time of 2 minutes. Detection was by Flame Ionization (Temperature 275°C, Fuel flow 30.0 ml/min, Oxidizer 400.0 ml/min). Instrument control and data collection and analysis was using the Millennium Chromatography Management System (Version 3.2, Waters Corporation, Milford, MA). Integration and quantification were performed automatically, but all analyses were subsequently examined manually to verify correct peak identification and acceptable signal to noise ratio before inclusion of the derived results in the study.

The ACTTAG line designated W000086431 was identified as having a high oil phenotype. Specifically, oil constituted 34.8% of seed mass (w/w) compared to an average oil content of 28.7% of other ACTTAG lines grown and analyzed in the same conditions (i.e. reference lines). Reanalysis of the same seed was performed in triplicate. Oil constituted 32.1% of seed mass, confirming an increase in oil content relative to reference.

The W000086431 line has three loci of T-DNA. One locus, was unlinked to the phenotype, and two closely linked loci were identified which cosegregated with the high oil phenotype. T2 individuals homozygous for the high oil loci produced seed with an oil content of 115.4% of the reference, T2 individuals hemizygous for the loci produced seed with an oil content of 118.4% of the reference. T2 individuals lacking the HIO-30 locus had oil contents of 105% of the reference. Because the homozygotes and hemizygotes for the high oil loci display a similar increase in oil content, it was determined that the high oil phenotype of W000068431 is dominant.

EXAMPLE 2

Characterization of the T-DNA Insertion in Plants Exhibiting the Altered Oil Content Phenotype.

We performed standard molecular analyses, essentially as described in patent application PCT WO0183697, to determine the site of the T-DNA insertion associated with

the altered oil content phenotype. Briefly, genomic DNA was extracted from plants exhibiting the altered oil content phenotype. PCR, using primers specific to the pSKI015 vector, confirmed the presence of the 35S enhancer in plants from line W000086431, and Southern blot analysis verified the genomic integration of the ACTTAG T-DNA.

5 Plasmid rescue and/or inverse PCR were used to recover genomic DNA flanking the T-DNA insertion, which was then subjected to sequence analysis using a basic BLASTN search and/or a search of the Arabidopsis Information Resource (TAIR) database (available at the arabidopsis.org website). Seeds from 18 T3 families descended from the mutant were harvested. The seed oil content of these families was determined as described in Example 1.
 10 The genotypes of these families with respect to a T-DNA insert were determined by T-DNA specific PCR using primers that are specific to one of the corresponding genomic region. The average oil content of T3 families containing the T-DNA insert at locus 2 and 3 was higher than those families lacking the insert at the corresponding loci. Therefore, we concluded that locus 2 and/or 3 is linked to the high oil phenotype. By contrast, the average
 15 oil content of T3 families containing the T-DNA insert at locus 1 was lower than /approximately the same as those families lacking the insert at the corresponding locus, and we concluded that locus 1 is not linked to the high oil phenotype.

Sequence analysis revealed that the start codon of the nucleotide sequence presented as SEQ ID NO: 1, which we designated HIO30.5, was approximately 1 kb 5' of the upstream
 20 border of the T-DNA insert at locus 3.

EXAMPLE 3

Analysis of *Arabidopsis* HIO30.5 Sequence

Sequence analyses were performed with BLAST (Altschul *et al.*, 1997, J. Mol. Biol.
 25 215:403-410), PFAM (Bateman *et al.*, 1999, Nucleic Acids Res 27:260-262), PSORT (Nakai K, and Horton P, 1999, Trends Biochem Sci 24:34-6), and/or CLUSTAL (Thompson JD *et al.*, 1994, Nucleic Acids Res 22:4673-4680).

BLASTN against GenBank sequences returned the candidate gene itself (plus its genomic DNA entry) and a putative Arabidopsis paralogue, At2g34900 (2 entries). The

At3g52280 candidate gene is not represented by a cDNA clone nor are there any Arabidopsis ESTs associated with this gene:

Sequences producing High-scoring Segment Pairs:

	Score	P(N)	N
gi 18409525 ref NM_115088.1 Arabidopsis thaliana chromos...	5550	5.6e-244	
gi 6434227 emb AL132972.1 ATT25B15 Arabidopsis thaliana D...	1155	1.0e-175	
gi 18403687 ref NM_129043.1 Arabidopsis thaliana chromos...	609	9.0e-19	1
gi 20197115 gb AC004238.3 Arabidopsis thaliana chromosom...	341	1.7e-05	2

BLASTP against amino acids in GenBank returned similar results, and identified 3 additional putative Arabidopsis homologues encoded by genes At2g34900 At5g10550 & At5g65630, having 39%, 13%, and 14% sequence identities, respectively, with SEQ ID NO:2. A predicted protein from rice (GI#7523514) was also identified, having 15% sequence identity with SEQ ID NO:2. The top 10 results are listed below and include human and fly genes encoding transcription factors that have been associated with controlling gene expression related to developmental patterns.

Sequences producing High-scoring Segment Pairs:

	Score	P(N)	N
gi 15231174 ref NP_190796.1 unknown protein; protein id:...	1570	1.1e-160	
gi 15226857 ref NP_181036.1 putative RING3 protein; prot...	606	1.5e-58	1
gi 15238195 ref NP_196617.1 bromodomain protein - like; ...	239	1.0e-26	2
gi 7523514 dbj BAA94242.1 Similar to Arabidopsis thalian...	278	3.4e-23	1
gi 15239091 ref NP_201366.1 putative protein; protein id:...	212	1.2e-21	2
gi 7657218 ref NP_055114.1 bromodomain-containing protei...	215	7.9e-21	2
gi 3184498 gb AAC27978.1 R31546_1 [Homo sapiens]	215	8.2e-21	2
gi 120558 sp P13709 FSH_DROME FEMALE STERILE HOMEOTIC PRO...	216	1.0e-20	2
gi 24640482 ref NP_511078.2 CG2252-PB [Drosophila melano...	216	1.0e-20	2
gi 1588281 prf 2208296A RING3 protein	255	1.2e-20	1

BLASTN of SEQ ID NO:1 against GenBank sequences identified the following Arabidopsis sequences with % identity to SEQ ID NO:1 indicated in parentheses: At2g34900 GI#15226857 (39%); At5g10550 GI#15238195 (13%) and At5g65630 GI#15239091 (14%).

BLASTN against ESTs returned the following ESTs from non-Arabidopsis plant species with % identity to SEQ ID NO:2 indicated in parentheses: potato (*Solanum tuberosum*) GI#s 13608255 (58%), and 21921887 (30%); cotton (*Gossypium* sp) GI#5050723 (61%); soy (*Glycine max*) GI#9835128 (30%) Brassica rapa GI#1019482 (51%); and corn (*Zea mays*) GI#24765979 (34%).

SEQ ID NOs 3 and 4 are sequences from tomato and wheat, respectively, that represent returned ESTs that were assembled into the least number of contigs. Where

contigs represent partial coding regions, the entire cDNA sequence can be determined by someone skilled in molecular biology techniques.

SEQ ID NO:3 is from tomato (*Lycopersicon esculentum*), and is a contig of the following sequences: GI#s 13778307, 10804738, 9504093, 6985312, 4381395. It has 58% identity with SEQ ID NO:1.

SEQ ID NO:4 is from wheat (*Triticum aestivum*) and is a contig of GI#s 21837181, 19950212, 20123686, 20547632, 19957908, and 20299669. It has 22% identity with SEQ ID NO:1.

PFam analysis detected a bromodomain (PF00439) from residues 87 to 200 of SEQ ID NO:2. In agreement with the predicted function as a transcription factor, PSORT2 analysis predicts that the At3g52280 gene product is nuclear (60%). This gene may play a regulatory role in the expression of genes involved in fatty acid metabolism or related pathways.

EXAMPLE 4

Confirmation of Phenotype/Genotype Association

RT-PCR analysis showed that the HIO30.5 gene was overexpressed in plants from the line displaying the HIO30.5 phenotype. Specifically, RNA was extracted from rosette leaves and/or siliques of plants exhibiting the HIO30.5 phenotype collected at a variety of developmental stages and pooled. RT-PCR was performed using primers specific to the sequence presented as SEQ ID NO:1, to other predicted genes in the vicinity of the T-DNA insertion, and to a constitutively expressed actin gene (positive control). The results showed that plants displaying the HIO30.5 phenotype over-expressed the mRNA for the HIO30.5 gene, indicating the enhanced expression of the HIO30.5 gene is correlated with the HIO30.5 phenotype.

The dominant inheritance pattern of the HIO30.5 phenotype is confirmed through genetic analysis. In general, genetic analysis involves the production and analysis of F1 hybrids. Typically, F1 crosses are carried out by collecting pollen from T2 plants, which is used to pollinate wild type plants. Such crosses are carried out by taking approximately 4 flowers from each selected individual plants, and using the T2 flower as the male pollen donor and flowers of the wild type plants as the female. 4-5 crosses are done for an

individual of interest. Seed formed from crosses of the same individual are pooled, planted and grown to maturity as F1 hybrids.

REFERENCES

- 5 Altschul, S.F. *et al.*, *J. Mol. Biol.* 215:403-410, 1990.
Altschul, S.F. *et al.*, *Nucleic Acids Res.* 25:3389-3402, 1997.
Ausubel FM *et al.* Current Protocols in Molecular Biology, John Wiley & Sons, New York, N.Y., 1993.
Baldwin D *et al.*, *Cur Opin Plant Biol.* 2(2):96-103, 1999.
- 10 Bateman *et al.*, 1999, *Nucleic Acids Res* 27:260-262 (website at pfam.wustl.edu).
Baulcombe D, *Arch Virol Suppl* 15:189-201, 1999.
Cannon *et al.*, *Plant Molec. Biol.* (1990) 15:39-47.
Ch'ng *et al.*, *Proc. Natl. Acad. Sci. USA* (1989) 86:10006-10010
Christensen S *et al.*, 9th International Conference on *Arabidopsis* Research. Univ. of
15 Wisconsin-Madison, June 24-28, 1998. Abstract 165.
Christou *et al.*, *Proc. Natl. Acad. Sci USA* (1989) 86:7500-7504.
Cough, SJ and Bent, AF, *the Plant Journal* 16(6): 735-743, 1998.
De Block *et al.*, *Plant Physiol.* (1989) 91:694-701.
Dieffenbach C and Dveksler G (Eds.) *PCR Primer: A Laboratory Manual*, Cold
20 Spring Harbor Laboratory Press, NY, 1989.
Everett *et al.*, *Bio/Technology* (1987) 5:1201
Feldmann *et al.*, *Science* 243: 1351-1354, 1989.
Focks N and Benning C, *Plant Physiol* 118:91-101, 1998.
Fridborg I *et al.*, *Plant Cell* 11: 1019-1032, 1999.
- 25 Geest AH and Hall TC, *Plant Mol Biol* 32(4):579-88, 1996.
Gelvin, S. B., Schilperoort, R. A., Varma, D. P. S., eds. *Plant Molecular Biology Manual* 1990.
Glick, BR and Thompson, JE, Eds. *METHODS IN PLANT MOLECULAR BIOLOGY AND BIOTECHNOLOGY*, p. 213-221, CRC Press, 1993.
- 30 Harlow E and Lane D, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, 1988, New York.
Harlow E and Lane D, *Using Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, 1999, New York
Hayashi H *et al.*, *Science* 258: 1350-1353, 1992.

- James DW and Dooner HK (1990) *Theor Appl Genet* 80, 241-245.
- Jensen, L.G., *et al.*, *Proc. Natl. Acad. Sci. USA* 93:3487-3491, 1996.
- Jones JD *et al.*, *Transgenic Res* 1:285-297 1992.
- Kardailsky I *et al.*, *Science* 286: 1962-1965, 1999.
- 5 Kline *et al.*, *Nature* (1987) 327:70.
- Kunkel TA *et al.*, *Methods Enzymol.* 204:125-39, 1991.
- Lemieux B, *et al.* 1990, *Theor Appl Genet* 80, 234-240.
- Nakamura Y *et al.*, 1999, *Nucleic Acids Res* 27:292.
- Napoli, *et al.*, *Plant Cell* 2:279-289, 1990.
- 10 Omirulleh *et al.*, *Plant Mol Biol.* 21(3):415-28, 1993.
- Sambrook *et al.* *Molecular Cloning: A Laboratory Manual* (Second Edition), Cold
Spring Harbor Press, Plainview, N.Y., 1989.
- Schaffer R, *et al.*, *Cell* 93: 1219-1229, 1998.
- Sheehy *et al.*, *Proc. Natl. Acad. Sci. USA* (1988) 85:8805-8809.
- 15 Smith, *et al.*, *Nature* 334:724-726, 1988.
- Smith *et al.*, *Mol. Gen. Genetics* (1990) 224:477-481.
- Thompson JD *et al.*, 1994, *Nucleic Acids Res* 22:4673-4680.
- van der Krol *et al.*, *Biotechniques* (1988) 6:958-976.
- van der Krol *et al.*, *The Plant Cell* (1990) 2:291-299.
- 20 Van Haaren MJJ *et al.*, *Plant Mol Bio* 21:625-640, 1993.
- Verdaguer B *et al.*, *Plant Mol Biol* 37:1055-1067, 1998.
- Waterhouse, *et al.*, *Proc. Natl. Acad. Sci. USA* 95:13959-13964, 1998.
- Weigel D, *et al.*, *Plant Physiology*, 122:1003-1013, 2000.
- Wilson K *et al.*, *Plant Cell* 8: 659-671, 1996.
- 25 Yadav NS *et al.* (1993) *Plant Physiol* 103, 467-476.

IT IS CLAIMED:

1. A transgenic plant comprising a plant transformation vector comprising a nucleotide sequence that encodes or is complementary to a sequence that encodes a HIO30.5 polypeptide, whereby the transgenic plant has a high oil phenotype.

2. The transgenic plant of Claim 1, which is selected from the group consisting of rapeseed, soy, corn, sunflower, cotton, cocoa, safflower, oil palm, coconut palm, flax, castor and peanut.

3. A plant part obtained from the plant according to Claim 1.

4. The plant part of Claim 3, which is a seed.

5. A method of producing oil comprising growing the transgenic plant of Claim 1 and recovering oil from said plant.

6. A method of producing a high oil phenotype in a plant, said method comprising:

a) introducing into progenitor cells of the plant a plant transformation vector comprising a nucleotide sequence that encodes or is complementary to a sequence that encodes a HIO30.5 polypeptide, and

b) growing the transformed progenitor cells to produce a transgenic plant, wherein said polynucleotide sequence is expressed, and said transgenic plant exhibits an altered oil content phenotype.

7. A plant obtained by a method of claim 6.

8. The plant of Claim 7, which is selected from the group consisting of rapeseed, soy, corn, sunflower, cotton, cocoa, safflower, oil palm, coconut palm, flax, castor and peanut.

9. The plant of claim 7, wherein the plant is selected from the group consisting of a plant grown from said progenitor cells, a plant that is the direct progeny of a plant grown

from said progenitor cells, and a plant that is the indirect progeny of a plant grown from said progenitor cells.

ABSTRACT

5 The present invention is directed to plants that display an altered oil content
phenotype due to altered expression of a HIO30.5 nucleic acid. The invention is further
directed to methods of generating plants with an altered oil content phenotype.

Nucleic Acid and Polypeptide Sequences

SEQ ID NO:1

>gi|18409525:1-1110 Arabidopsis thaliana chromosome 3 CHR3v07142002 genomic sequence

ATGGCTGACTCAGTGCCAGGTCATGTCGCCGGCGGAGGTTTACAGGGTTTCTCTGTCGACGCCGAGTGCA
TCAAGCAGCGTGTGTCAGCAGGTTTACAGTGGGTTGATTCGCTTGAGCATAAACTGAAAGAAGTGGAGGA
GTTTTACTCGAGCATTTGGTGTCTCAAACTCTGGCTCTATTGGTAAAGATACAGAAAAAGGGAGACATGTT
GTGGGGATTAGGAAGATTCAACAAGAGGCTGCGCGTAGAGAAGCTGTTGCTGCTAAAAGAATGCAGGACC
TTATGCGTCAATTTGGAACAATCTTTCTGTCAGATAACTCAGCATAAGTGCCTTGGCCGTTTATGCATCC
TGTCATGTTGAAGGTCTTGGTTTGCATGACTACTTTGAGGTTATTGACAAGCCTATGGACTTCAGCAG
ATAAAAAATCAAAATGGAGGCTAAGGATGGTACCGGTACAAACATGTCATGCAAAATTTATGCTGATATGC
GGCTAGTGTTTGAGAACGCAATGAATTATAATGAAGAAACAAGTGATGTTTACTCTATGGCGAAAAAGTT
ACTGGAAGAGTTTGAAGGAGAGTGGGCACACTTTCTTCCAAAGTTCAAGAAGAGGAGAAAAATACGAGAG
GAAGAGGAGAGCAAGCAGCAAGGAGGCGCTGCTAGCAAAAGAAGCATCTCATATCAAAACAACTAGAG
AATTGGGGAAATGAGATTGTCATGCTAATGACGAGCTGGAGAAGCTAATGCGTAAAGTTGTTGGAAAGATG
CAGGAAAAATCACAATTGAGGAGAGCGTAATATTGGATTGGCAATTGTTAAACTGTCACCGGATGATCTA
CAGAAAGTGTGGGTATAGTTGCTCAGGCTAACCGAGCTTTCAACCTAGAGCAGAGGAAGTTAGTATTG
AGATGGACATCTGGACGAACCACTATGGAGGCTAAAGTCTTTGTTGAAGGATGCGTTGGATAATGC
AATGAAGAAGAAGAAGAGGAGACAAAGACTAGAGAATTGAGTGGAGCACAAAAGAAAGAGGTTTGC
AAGAAACGAAATGCGCATACAAAGTTAGCTGAGAGGAAAAAGCGGTACGCATATGA

SEQ ID NO:2

>gi|15231174|ref|NP_190796.1| unknown protein; protein id: At3g52280.1 [Arabidopsis thaliana]

MADSVPGHVAGGLQGFVDAECIKQRVDEVLLQWVDSLEHKLKEVEEFYSSIGVSNSSIGKDTKEGRHV
VGIRKIQEEARREAVAARKMQDLMRQFGTIFRQITQHKCAWPFMHPVNVVEGLGLHDYFEVIDKPMDFST
IKNQMEAKDGTGYKHVMQIYADMLRVFENAMNYNEETSDVYSMAKKLLEKFEKWAHFLPKVQBEKIRE
BEEKQAAKEALLAKEASHIKTTRELGNIEICHANDELEKLMRKVVERCRKITEEKRNIGLALLKLSPPDDL
QKVLGIVAQANPSFQPRAEVSIEMDILDEPTLWRLKFFVKDALDNAMKKKKEETKTRELGAQKKEVS
KKRNATTKLAERKTKRSRI

SEQ ID NO:3

GCCTTCATCGGCGGCACAACCTGTTGCTTATCTCCGATCTACTCCGCGAACGGCGGAAAT
TCGAGCTATGGAGAACTTAAACCGCTTAGTTCTTGACATCCCTATACAAGAGCAGAAAAGA
CAATGCTACTGCTTCTGAAAATTTCCGTCGAAAGTGTGATGAAATGGTTGGGAAGGTTGA
TCAGATTGAGCAGAGACTGAATGAGGTTGAGCATTCTATTTCCAATACTAGTAAAAACA
ATCAAAACACGCCAAGAGGTTGGCTCTATTCTGAAAGACAAAGAGAAACAGATGCTAGCTT
TAGAAGGGCGGCAGCAGGATGCATCAGTAGAGAAGCAGCTGGTTCCAGGAGAAATGCAAGA
ACTTATGCGCCAAATTTGGTACTATATTACGTGAGATCACTCAGCACAAGTGGGCGAAGC
TTTTATGGAACCTGTGGATGTCAAGGGTCTTGGATTACACGATTATTTTGAGGTTATTGA
GAAGCCTATGGACTTCAGTACTATCAAAACAAAGATGGAAGCAAGGATGGTTCTGGCTA
TAAGCATGTCCGAGAGATGTGCTGACGTGAGGCTAATATTTAAGAATGCGATGAAATA
CAACGAGGAAAGGGATGACGTTTATGTAATGGCGAAAACCTTGTGGGAAAATTCGAGGA
GAAATGGTTGCAGCTTTTGCCCAAAGTTGATGAAGAGGAAAAACGGCGAAAGGAAGAGGA
AGCAGAAGCCCAACAGGATATGCAGCTCGCTCAAGAGGCTGCTCATGCAAAAATGGCAAA
AGATTTAACTATTGAGCTTGATGAGGTGACATGCAACTGGAAGAACTCAGGGAATTTGGT
GCTTCAGAAATGCAGAAAGATTTCAACGGAAGACAGGAAACAACTTGGGAATGCGCTCAC
TAAGCTGTCTCCTGACGATCTTAATAAAGCACTATTGATTGTGCGACAGAATGATCTTAC
TTTTTC

SEQ ID NO:4

CGAGGGGCGGCAATGGAGGGAGCGTGGCCCCATCCGCGCGCGATGGAGCCGACCACCGCC
GCGGATGGGCCCTCAAGTGTGCGAGGTGAACCTGTTCCGCGCGCAGGTCGACGACCTCCTC
TCCAAGACCGACGTGCTGGAGAAGAGGGTGAACGAGGTGGTAGGGTTCTACAATTTCAAAG
AAGCACAGCAGTGGAGGCGCAAGGCTGGCGCGAGGTACGCGGAAAACGGTGCCAGGGAC
AGCCACGGCAATGGGATGCCGACCTCATGCGCCAGTTTGCCGCTATCATTGCGCAGATT
ACATCTCATGAATGGGCACAGCGGTTTTTGCAACCGGTAGACGTCGTAGGTCTTCAACTT
GATGACTATCACCAGATTATAACAAAACCTATGGATTCTCGACCATCCGAAACAAAATG
GAAGGGAAGGAAAGTACCACATATAACAGTGTCCGAGAAATATATTCTGATGTAGATTA
GTTTTTACCAATGCAATGAAGTACAATGTTGAAGGTACCCCTGTTAACATAATGGCCAAG
TTCTACTCGAGAGATTGAGGAGAAATGGCTTCACCTTCTCCCTGAAGTTGAGAACGAG
GAAAGGGAACGGGAGGAACCAATGATGCTCCAACCATAAAGCATTTCTCCGGAAGCTGCA
ATTGCAAAATTAGCTGAAGATACTGGTAATGAGCTGAATGAGATTAATAAGCAGCTGGAG
GAGCTCCAGAAAATGGTGGTTTCAGAGATGCAGGAAAATGACGACGGACGAGAGAGAAAA
CTCGGCGCAGGCTTTTGGCAATTGTCTCCAGAAGATCTTAACAAGGCGCTAGAGTTGGTC
GCGCAAGACAATCTAGCTTCCAAACTACAGCTGAAGAAGTGGACCTTGACATGGATGCT
CAGAGCGAGACGACCTCTGGAGGCTGAAGTTCTTTGTGAGGGAAGCATTGGAACAACAG
CGAATGTAGGCGTAGTGGCTGTGGCAAGATCGATGAAAAACGAAGAGGAGCCGCGAC

5

ATGTACAATGCTCTAGCCAAGACCGTTTCGAAACGGATTAAGAGATAACCTTAGCGGCTC
GGCCTTCTGATGGTTCATGGTATTTCTGTAGAAAATAGGGCCATGTTCTCCCTTGTGTG
AAGACATGTTGTATATATACGAAGATGAATCTGCACTAAGATATGCTTGGTCGCACCCGG
TCTCATGTGGCATGGGTAGCATGCTATTGGTAGCTATGTAAAAATAAGAGCAATGTTGAA
CACCTCTGAGAAACAAGAATTATTCCTTGA